



## Paternal Postnatal Depression and its Risk factors: A systematic review of cross-sectional studies

Dhanpal HN, Principal, Victory College of Nursing, Bangalore

[dhanpal2000@gmail.com](mailto:dhanpal2000@gmail.com)

### Abstract

**Background and Aim:** Postnatal depression (PND) can affect both genders, but the common misconception is that it only affects mothers. Increasing literature reports that 10% of the fathers experience PND after childbirth. This systematic review aims to ascertain the prevalence of PND in fathers and its risk factors.

**Materials and Methods:** The electronic databases PubMed, BASE, DOAJ, Research Gate Semantic Scholar, and BioMed Central were searched for related open access articles published between 2010 to December 2021. Finally, 12 articles met inclusion criteria.

**Results:** The findings reported that the prevalence of PND in fathers ranged from 5.7% to 59.8%. Partner's depression, lack of social support, poor marital relationship, low income, and low education were all shown to raise the risk of PND in fathers.

**Conclusion:** PND in fathers is a serious concern. Early identification and treatment decrease the detrimental impact on mother and child while further improving quality of life.

**Keywords:** Father, Postnatal Depression, Postpartum Depression, Edinburgh Postnatal Depression Scale, Prevalence, Risk factors

### Background

Postnatal depression (PND) or Postpartum depression (PPD) is a non-psychotic depressive disorder that occurs within the first year after childbirth<sup>1-2</sup>. PND can affect both parents but it has been associated primarily with mothers<sup>3-4</sup>. However, increasing literature shows that PND is not uncommon in fathers and 10% or 1 in 10 fathers around the world experience depression after childbirth<sup>5</sup>. The prevalence of PND in fathers during the first year after childbirth has been found to range from 4% to 25%, with a 50% rise when the mother is also depressed<sup>4-6</sup>. A more recent meta-estimate recorded for PND in fathers within the postpartum period was 8.4%<sup>7</sup>. This rate is higher than the depression rate (4.7%) in the general male adult population<sup>8</sup>.

Several factors that precipitate PND in fathers have been reported by researches including partner's depression, previous history of depression, unemployment, low education, poor marital relationship, lack of social support, and an unplanned pregnancy<sup>9-11</sup>. The partner's depression during the postnatal period was found to be the best predictor of PND in fathers<sup>11</sup>.



PND has a detrimental influence on fathers' health and well-being, as well as that of their families. It involves failure to fulfill obligations at home and at work, a loss of interest, fatigue, stress, and a higher risk of suicide in fathers<sup>12-14</sup>. Complications in marital relationships and the development of behavioral and emotional problems in their children are some of the negative impacts of PND in fathers on their families<sup>15-17</sup>.

PND in fathers is a clinically significant problem with higher community-based care costs<sup>18</sup>. Despite this, PND in fathers is under-screened, under-diagnosed, and under-treated<sup>19-20</sup>. Various scholars and countries have investigated PND in fathers, but it is still in its early stages in India. The goal of this review is to raise the understanding regarding PND in fathers and its associated risk factors. Thus this review aims to ascertain the prevalence of PND in fathers and to determine its risk factors.

## **Methodology**

The systematic review reporting follows Preferred Reporting Items for Systematic reviews and Meta-analyses (PRISMA) guidelines.

### **Search strategy**

PubMed, BASE, DOAJ, Research Gate Semantic Scholar, and BioMed Central electronic databases were searched for articles published between 2010 to December 2021 by using the following terms with Boolean operators ('OR'/'AND'): "father" OR "paternal" AND "Postnatal depression" OR "postpartum depression".

### **Inclusion and exclusion criteria**

Articles were chosen for inclusion based on the following criteria: 1) Journal articles that examined the prevalence of PND in fathers and its associated factors, 2) Cross-sectional research design, 3) Published in the English language, 4) Published between 2010 to December 2020, 5) Original research articles and 6) Full free and open access articles. Exclusion criteria were: 1) Journal articles abstracts, Review articles, Commentaries, Conference Reports, and Thesis, 2) Interventional articles, 3) Duplication, and 4) Articles not relevant to the study.

### **Data Extraction**

The data was extracted by the author with the help of experts. The data extracted from the eligible studies were: year of citation, author, country, research design, sample size, response rate, recruitment setting, assessment points, assessment instrument, cut-off score, and prevalence.



## Methodological quality Assessment

Two independent authors used the Mirza and Jenkins checklist to determine the methodological quality of the included studies<sup>21-22</sup>. The critical appraisal checklist included the following criteria: 1) clear study objectives, 2) adequate sample size (or justification), 3) representative sample (with justification), 4) clear inclusion and exclusion criteria, 5) depression measure used is reliable and valid, 6) reported response rate and/or losses explained, 7) adequate description of data, 8) appropriate statistical analyses and with additional criterion 9) appropriate informed consent<sup>23</sup>. Each criterion of the checklist is answered by 'yes' (1 point) or 'no' (no point). Based on the points obtained by the studies were graded between 1 to 9. Regardless of their quality, all of the studies were included.

## Results

### Search results and inclusion of articles

A total of 1085 articles were found through a database search. 998 articles were excluded after an initial screening for a variety of reasons (Figure 1). A total of 43 full free text articles were chosen and screened against inclusion criteria. Finally, 12 articles met the inclusion criteria.




Figure 1: PRISMA: Literature Search Flow Chart



### Study characteristics

All the 12 cross-sectional studies selected<sup>24-35</sup>, investigated the prevalence of postnatal depression and its associated factors. Included studies were conducted in Iran (3)<sup>24, 26, 29</sup>, Japan (1)<sup>25</sup>, Italy (1)<sup>27</sup>, Sweden (1)<sup>28</sup>, Ethiopia (1)<sup>34</sup>, Saudi Arabia (3)<sup>32, 33, 35</sup>, Ireland (1)<sup>30</sup>, and Chile (1)<sup>31</sup>. The key features of the studies included are briefly listed in Table 1.

### Quality assessment

According to the previously stated criterion, the quality of the included studies varied from 6 to 9 out of a possible maximum of 9, suggesting that these studies were of high methodological quality in general. Table 2 summarizes the findings of this assessment.

### Description of study subjects

Only fathers were the research subjects in eight of the twelve studies<sup>24,26,28,30,32-35</sup> (two of which included the first-time fathers<sup>24,26</sup>), while couples were the study subjects in the remaining studies<sup>25,27,29,31</sup> (one of which included first-time parents<sup>27</sup>).

### Sample size and Sampling technique

In studies with only fathers as a sample, the sample size ranged from 100 to 8011<sup>24,26,28, 30,32-35</sup> and in studies with couples as a sample, the sample size ranged from 75 to 2032<sup>25,27,29,31</sup>. Five of the 12 studies used probability sampling techniques<sup>24,28,30,32,34</sup>, two used non-probability sampling techniques<sup>26,33</sup>, and the others didn't mention.

### Study setting and time of recruitment of participants

The studies included in this review varied in terms of the time and setting in which participants were recruited. The participants of the studies were recruited at different time points from childbirth to 12 months of postpartum. Participants were recruited in five studies within 3 months of postpartum<sup>24,26,27,29,35</sup>, two studies between 3 and 6 months<sup>25,28</sup>, and two studies between birth and 12 months<sup>30,33</sup>. Of the remaining studies, one study enrolled participants after two months of childbirth<sup>31</sup>, one study from childbirth to 6 months<sup>32</sup>, and one study after 4 weeks of postpartum<sup>34</sup>. Three studies recruited a sample from health centre<sup>24,28,33</sup>, and 3 studies from hospital<sup>26,31,35</sup>. Out of the remaining six studies, one study recruited a sample from population registry<sup>28</sup>, one study directly from the community<sup>25</sup>, one study from private pediatric ambulatories<sup>27</sup>, one study from primary health care centre<sup>30</sup>, one from tertiary care<sup>31</sup> and one study recruited its sample from health care centre<sup>34</sup>.

### Tool used to measure paternal postnatal depression.

The Edinburgh Postnatal Depression Scale (EPDS) was used in all of the studies to screen fathers for PND. However, the cut-off score for the PND screening varied. For positive PND screening, four studies used an EPDS cut-off score of  $\geq 12$ <sup>24,26,27,29</sup>, three studies used a cut-off



**Table 1: Methodological characteristics of studies included in the systematic review**

Year of citation/ Authors / Country	Study Design	Sample size	Response Rate	Recruitment setting	Assessment time	Assessment Instruments	Cut-off score	Prevalence
2014 Kamalifard et al. Iran	Cross-sectional	205 New fathers	97%	7 Health centers	-At 6 <sup>th</sup> to 12 <sup>th</sup> week postpartum	EPDS	≥ 12	11.7%
2015 Nishimura et al. Japan	Cross-sectional	2032 couples.	39.7% couples	4 Wards of Kobe city	-At 4 months of postpartum	EPDS	≥ 8	13.6. %
2015 Ahmadi Z et al. Iran	Cross-sectional	328 New fathers	100%	Hospital	-At 8 <sup>th</sup> week of postpartum	EPDS	> 12	59.8%
2015 Epifanio MS et al. Italy	Cross-sectional	75 couples (first-time parents)	70.6%	Private pediatric ambulatories	-At first month of child's birth	EPDS	> 12	5.7%
2018 Carlberg M et al. Sweden	Cross-sectional	8,011 fathers	46%	Population registry.	-3 to 6 months postpartum	EPDS GMDS	≥ 10 ≥ 12 ≥ 13	13.3% 8.1% 8.6%
2018 Kamalifard M et al. Iran	Cross-sectional	205 couples	90%	7 Health centers	- At 6 <sup>th</sup> to 12 <sup>th</sup> weeks after childbirth.	EPDS	≥ 12	11.7%
2018 Philpott LF and Corcoran P Ireland	Cross-sectional	100 fathers	77%	Primary care center	-Birth to a child in the last 12 months	EPDS	≥ 9 ≥ 12	28% 12%



(Table 1 continued)

Year of citation/ Authors / Country	Study Design	Sample size	Response Rate	Recruitment setting	Assessment time	Assessment Instruments	Cut-off score	Prevalence
2018 Francisca Perez C et al. Chile	Cross-sectional	382 couples	33.5%	Hospital	-2 months after child birth	EPDS BDI	≥ 10 13/14	18.5 10.5
2019 Shaheen et al. Saudi- Arabia	Cross-sectional	347 fathers	83.57%	Tertiary care	-From child's birth to 6 months postpartum	EPDS DSM-5 criteria	8/9	16.6%
2019 AlHaisoni MS Saudi Arabia	Cross-sectional	226 fathers	85.2%	Health care Center	-Up to 12 months of postpartum	EPDS	≥ 9	32.7%
2020 Markos and Arba Ethiopia	Cross-sectional	423 fathers	97%	25 public health centers	≥ 4 weeks of postpartum	EPDS	≥10	17%
2020 Alghamdi et al. Saudi Arabia	Cross-sectional	182 fathers	82.41%	Hospital	-Between 4 and 8 weeks of postpartum	EPDS	>10	27.3%

**EPDS** - Edinburgh Postnatal Depression Scale; **BDI** - Beck Depression Inventory; **GMDS** - Gotland Male Depression Scale; **DSM-5** - Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition

score of  $\geq 10^{31,34,35}$ , two studies used a cut-off score of  $\geq 9^{32,33}$ , and one study used an EPDS cut-off score of  $\geq 8^{25}$ . The other two researches used two cut-off scores: one used a cut-off score of  $\geq 10$  and  $\geq 12$  for positive screening of minor and major depression<sup>28</sup>, and the other used a cut-off score of  $\geq 9$  and  $\geq 12$  for positive screening of minor and major depression<sup>30</sup>. The Beck Depression Inventory (BDI) with a cut-off score of 13/14<sup>31</sup> and the Gotland Male Depression Scale (GMDS) with a cut-off score of  $\geq 13^{28}$  were also used as screening measures. To compare the screening findings, two studies used the BDI and GMD with EPDS<sup>28,31</sup>. After screening with EPDS, one study used the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) to determine a PND diagnosis<sup>32</sup>.

### Prevalence of PND

According to studies, the overall prevalence rate of PND in fathers ranges between 5.7% and 59.8%<sup>24,35</sup>. The lowest prevalence was reported in Italy<sup>27</sup> and the highest was in Iran<sup>26</sup>. PND was observed in 5.7% to 59.8% of fathers between the childbirth to 3 months<sup>24,26,27,29,35</sup>, 13.3% to 13.6% between three and six months<sup>25,28</sup>, and 28% to 32.7% between childbirth and 12 months of postpartum<sup>30,33</sup>. The prevalence ranged from 5.7% to 59.8% when using an EPDS cutoff score of  $\geq 12^{24,26-29}$ , 13.3% to 27.3% when using a cutoff score of  $\geq 10^{28,31,34,35}$ , and 16.6% to 32.7% when using a cutoff score of  $\geq 9^{32,33}$ . When two cutoff scores were used, the prevalence ranged from 8.1% to 13.3% in one study<sup>28</sup> and from 12% to 28% in another<sup>30</sup>.

Study	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>	C <sub>4</sub>	C <sub>5</sub>	C <sub>6</sub>	C <sub>7</sub>	C <sub>8</sub>	C <sub>9</sub>	Total score
Kamalifard M et al. <sup>24</sup>	1	1	1	1	1	1	1	1	1	9
Nishimura A et al. <sup>25</sup>	1	1	1	0	1	1	1	1	1	8
Ahmadi Z et al. <sup>26</sup>	1	1	0	0	1	0	1	1	1	6
Epifanio, M. S et al. <sup>27</sup>	1	0	0	0	1	1	1	1	1	7
Carlberg M et al. <sup>28</sup>	1	1	1	0	1	1	1	1	1	8
Kamalifard M, et al. <sup>29</sup>	1	1	1	1	1	1	1	1	1	9
Philpott LF & Corcoran P <sup>30</sup>	1	0	1	0	1	0	1	1	1	7
Francisca Pérez C et al. <sup>31</sup>	1	1	1	1	1	1	1	1	1	9
Shaheen NA et al. <sup>32</sup>	1	0	1	1	1	1	1	1	1	8
AlHaisoni MS <sup>33</sup>	1	1	1	1	1	0	1	1	1	8
Markos M & Arba A <sup>34</sup>	1	1	1	1	1	1	1	1	1	9
Alghamdi W A et al. <sup>35</sup>	1	0	1	1	1	1	1	1	1	8

**Table 2: Methodological quality assessment of studies included in the systematic review**



**C1**-Clear study objectives; **C2**-Adequate sample size or justification; **C3**-Representative Sample (with justification); **C4**- Clear inclusion and exclusion criteria; **C5**-Depression measure used is reliable & valid; **C6**-Reported response rate and/or losses explained; **C7**-Adequate description of data; **C8**-Appropriate statistical analyses; **C9**-Appropriate informed consent.

### Risk factors

Out of 12 studies, 11<sup>24-32,34,35</sup> studies reported risk factors for PND in fathers. Partner's depression<sup>25,28,29,31</sup>, lack of social support<sup>24,30,34</sup>, poor marital partnership satisfaction<sup>25,34,35</sup>, low income<sup>28,30,34</sup>, low education<sup>26,28</sup>, perceived stress<sup>24,35</sup>, and infant sleep problems<sup>30,34</sup> were the most frequently identified risk factors. Other factors reported were history of infertility treatment, economic anxiety, the experience of visiting medical institutions due to mental health problems<sup>25</sup>, unemployment<sup>26</sup>, maternal distress<sup>27</sup>, family livelihood situation<sup>29</sup>, history of depression, no paternity leave<sup>30</sup>, feeling isolated and disconnected from partner<sup>32</sup>, substance use, unplanned pregnancy<sup>34</sup>, family and work-related related problems, trouble sleeping, and low self-esteem<sup>35</sup>.

### Discussion

In the present study, the prevalence of PND in fathers was found to range from 5.7% to 59.8%. This wide variation may be attributed to varying cut-off points, sample sizes, cultural contexts, and assessment time<sup>5,7</sup>. The other reason for the wide variation in the prevalence rates was the use of self reporting measures. It has been observed that self-reporting tools provide high prevalence estimates than the diagnostic tools and interview<sup>37,38</sup>.

The most frequently found risk factors in the current review were partner's depression, lack of social support, poor marital partnership satisfaction, low income, and low education, which is consistent with previous findings<sup>9-11</sup>. Other common risk factors identified by previous researches such as unemployment and previous history of depression<sup>39,40</sup> were not reported by the majority of the studies (10/12) under the current review.

### Limitations

The review's limitations include the small number of identified studies, cross-sectional prevalence studies, and the variability in research methodology.

### Conclusion

According to a systematic review undertaken in India, 22% of mothers experience PND<sup>41</sup>. Mothers' PND is a strong indicator of PND in their partners. Furthermore, when their partners





were depressed in the postpartum period, PND in fathers ranged from 24% to 50%<sup>4</sup>. Nonetheless, PND in mothers gains more emphasis. PND in fathers is a serious concern and must be regarded as a public health issue. The mental health of fathers should be included in postpartum mental health assessments as early diagnosis and treatment decrease the detrimental impact on mother and child while further improving quality of life.

## Financial support and sponsorship

Nil

## Conflict of Interest

There are no conflicts of Interest

## Acknowledgement

I thank Rahul Shil, and Shivaleela P Upashe for performing data extraction quality assessment and providing valuable suggestions in inclusion of relevant articles.

## References

1. Massoudi P, 2013. Depression and distress in Swedish fathers in the postnatal period. Available at: [https://gupea.ub.gu.se/bitstream/2077/32509/1/gupea\\_2077\\_32509\\_1.pdf](https://gupea.ub.gu.se/bitstream/2077/32509/1/gupea_2077_32509_1.pdf) (Accessed 12 December 2016)
2. Matthey S, Barnett B, Ungerer J, Waters B. Paternal and maternal depressed mood during the transition to parenthood. *J Affect Disord.* 2000;60(2):75-85. doi: 10.1016/s0165-0327(99)00159-7, PMID 10967366.
3. O'Hara MW, Swain AM. Rates and risk of postpartum depression: A meta-analysis. *Int Rev Psychiatry.* 1996;8(1):37-54. doi: 10.3109/09540269609037816.
4. Goodman JH. Postpartum depression beyond the early postpartum period. *J Obstet Gynecol Neonatal Nurs.* 2004;33(4):410-20. doi: 10.1177/0884217504266915, PMID 15346666.
5. Paulson JF, Bazemore SD. Prenatal and postpartum depression in fathers and its association with maternal depression: A meta-analysis. *JAMA.* 2010;303(19):1961-9. doi: 10.1001/jama.2010.605, PMID 20483973.
6. Earls MF, Yogman MW, Mattson G, Rafferty J. Incorporating recognition and management of perinatal depression into pediatric practice. *Pediatrics.* 2019;143(1). doi: 10.1542/peds.2018-3259, PMID 30559120.
7. Cameron EE, Sedov ID, Tomfohr-Madsen LM. Prevalence of paternal depression in pregnancy and the postpartum: an updated meta-analysis. *J Affect Disord.* 2016;206:189-203. doi: 10.1016/j.jad.2016.07.044, PMID 27475890.



8. National Institute of Mental Health. Major depression among adults; 2015. Available at: <https://www.nimh.nih.gov/health/statistics/prevalence/majordepressionamongadults.shtml> . (Accessed 12 December 2016).
9. Escriba-Agüir V, Artazcoz L. Gender differences in postpartum depression. A longitudinal cohort study. *J Epidemiol Commun Health*. 2011;65(4):320-6. doi: 10.1136/jech.2008.085894, PMID 20515899.
10. Gao LL, Chan SW, Mao Q. Depression, perceived stress, and social support among first-time Chinese mothers and fathers in the postpartum period. *Res Nurs Health*. 2009;32(1):50-8. doi: 10.1002/nur.20306, PMID 18980171.
11. Paulson JF, Bazemore SD, Goodman JH, Leiferman JA. The course and interrelationship of maternal and paternal perinatal depression. *Arch Womens Ment Health*. 2016;19(4):655-63. doi: 10.1007/s00737-016-0598-4, PMID 26790687.
12. Veskrna L. Peripartum depression—does it occur in fathers and does it matter? *J Mens Health*. 2010;7(4):420-30. doi: 10.1016/j.jomh.2010.10.004.
13. Quevedo L, da Silva RA, Coelho F, Pinheiro KA, Horta BL, Kapczynski F, Pinheiro RT. Risk of suicide and mixed episode in men in the postpartum period. *J Affect Disord*. 2011;132(1-2):243-6. doi: 10.1016/j.jad.2011.01.004, PMID 21277023.
14. Kamalifard M, Hasanpoor S, Babapour Kheiroddin J, Panahi S, Bayati Payan S. Relationship between fathers' depression and perceived social support and stress in postpartum period. *J Caring Sci*. 2014;3(1):57-66. doi: 10.5681/jcs.2014.007, PMID 25276749.
15. Kerstis B, Berglund A, Engström G, Edlund B, Sylvén S, Aarts C. Depressive symptoms postpartum among parents are associated with marital separation: A Swedish cohort study. *Scand J Public Health*. 2014;42(7):660-8. doi: 10.1177/1403494814542262, PMID 25053465.
16. Ramchandani P, Stein A, Evans J, O'Connor TG, ALSPAC study team. Paternal depression in the postnatal period and child development: A prospective population study. *Lancet*. 2005;365(9478):2201-5. doi: 10.1016/S0140-6736(05)66778-5, PMID 15978928.
17. Paulson JF, Keefe HA, Leiferman JA. Early parental depression and child language development. *J Child Psychol Psychiatry*. 2009;50(3):254-62. doi: 10.1111/j.1469-7610.2008.01973.x, PMID 19175819.
18. Edoka IP, Petrou S, Ramchandani PG. Healthcare costs of paternal depression in the postnatal period. *J Affect Disord*. 2011;133(1-2):356-60. doi: 10.1016/j.jad.2011.04.005, PMID 21561664.
19. Brockington I. Postpartum psychiatric disorders. *Lancet*. 2004;363(9405):303-10. doi: 10.1016/S0140-6736(03)15390-1.
20. Musser AK, Ahmed AH, Foli KJ, Coddington JA. Paternal postpartum depression: what health care providers should know. *J Pediatr Health Care*. 2013;27(6):479-85. doi: 10.1016/j.pedhc.2012.10.001, PMID 23182851.
21. Mirza I, Jenkins R. Risk factors, prevalence, and treatment of anxiety and depressive disorders in Pakistan: systematic review. *BMJ*. 2004;328(7443):794. doi: 10.1136/bmj.328.7443.794, PMID 15070634.



22. Sawyer A, Ayers S, Smith H. Pre- and postnatal psychological wellbeing in Africa: A systematic review. *J Affect Disord.* 2010;123(1-3):17-29. doi: 10.1016/j.jad.2009.06.027, PMID 19635636.
23. Fisher J, Cabral de Mello M, Patel V, Rahman A, Tran T, Holton S, Holmes W. Prevalence and determinants of common perinatal mental disorders in women in low- and lower-middle-income countries: A systematic review. *Bull World Health Organ.* 2012;90(2):139G-49G. doi: 10.2471/BLT.11.091850, PMID 22423165.
24. Kamalifard M, Hasanpoor S, Babapour Kheiroddin J, Panahi S, Bayati Payan S. Relationship between fathers' depression and perceived social support and stress in postpartum period. *J Caring Sci.* 2014;3(1):57-66. doi: 10.5681/jcs.2014.007, PMID 25276749.
25. Nishimura A, Fujita Y, Katsuta M, Ishihara A, Ohashi K. Paternal postnatal depression in Japan: an investigation of correlated factors including relationship with a partner. *BMC Preg Childbirth.* 2015;15:128. doi: 10.1186/s12884-015-0552-x, PMID 26026489.
26. Ahmadi Z, Jan-Mohammadi S, Hajkazemi E. Prevalence of parental postnatal depression in fathers and its relationship with demographic characteristics. *JCCNC.* 2015;1(3):159-64.
27. Epifanio MS, Genna V, De Luca C, Roccella M, La Grutta S. Paternal and maternal transition to parenthood: the risk of postpartum depression and parenting stress. *Pediatr Rep.* 2015;7(2):5872. doi: 10.4081/pr.2015.5872, PMID 26266033.
28. Carlberg M, Edhborg M, Lindberg L. Paternal perinatal depression assessed by the Edinburgh Postnatal Depression Scale and the Gotland male depression scale: prevalence and possible risk factors. *Am J Mens Health.* 2018;12(4):720-9. doi: 10.1177/1557988317749071, PMID 29350097.
29. Kamalifard M, Bayati Payan S, Panahi S, Hasanpoor Sh, Babapour Kheiroddin J. Paternal postpartum depression and its relationship with maternal postpartum depression. *J Holist Nurs Midwifery.* 2018;28(2):115-20. doi: 10.29252/HNMJ.28.2.115.
30. Philpott LF, Corcoran P. Paternal postnatal depression in Ireland: prevalence and associated factors. *Midwifery.* 2018;56:121-7. doi: 10.1016/j.midw.2017.10.009, PMID 29096280.
31. Francisca Pérez C, Catalán A, Morales A, Quinlan A, Riquelme D, Navarro J, Valdés C. Assessment of postpartum depression in a group of Chilean parents. *J Mens Health.* 2018;14(2):e56-64. doi: 10.22374/1875-6859.14.2.7.
32. Shaheen NA, AlAtiq Y, Thomas A, Alanazi HA, AlZahrani ZE, Younis SAR, Hussein MA. Paternal postnatal depression among fathers of newborn in Saudi Arabia. *Am J Mens Health.* 2019;13(1):1557988319831219. doi: 10.1177/1557988319831219, PMID 30776946.
33. AlHaisoni MS, Ayman Afifi M. Prevalence of paternal postpartum depression among fathers visiting Well-Baby Clinic at Prince Sultan military medical city 2018. *IJMSCR.* 2019;2(4):486-97.
34. Markos M, Arba A. Prevalence and predictors of postpartum depression among Male Partners who came to postnatal Follow-Up Clinic with their partner in selected Public Health Centers of Wolaita Zone, Ethiopia, 2019. *Neuropsychiatr Dis Treat.* 2020;16:2307-16. doi: 10.2147/NDT.S273045, PMID 33116529.



35. Alghamdi WA, Alyahyawi HY, Shaaban SS, Alharbi RS, Yaghmoor BE, Alnajjar SF, Alshamrani HM. Prevalence and correlates of paternal postpartum depression in Saudi Arabia. *J Mens Health*. 2020;16(1):e41-52. doi: 10.15586/jomh.v16i1.199.
36. Ramchandani PG, Psychogiou L, Vlachos H, Iles J, Sethna V, Netsi E, Lodder A. Paternal depression: an examination of its links with father, child, and family functioning in the postnatal period. *Depress Anxiety*. 2011;28(6):471-7. doi: 10.1002/da.20814, PMID 21506206.
37. Ji S, Long Q, Newport DJ, Na H, Knight B, Zach EB, Morris NJ, Kutner M, Stowe ZN. Validity of depression rating scales during pregnancy and the postpartum period: impact of trimester and parity. *J Psychiatr Res*. 2011;45(2):213-9. doi: 10.1016/j.jpsychires.2010.05.017, PMID 20542520.
38. O'Hara MW, Swain AM. Rates and risk of postpartum depression-A meta-analysis. *Int Rev Psychiatry*. 1996;8(1):37-54. doi: 10.3109/09540269609037816.
39. Ramchandani PG, Stein A, O'Connor TG, Heron J, Murray L, Evans J. Depression in men in the postnatal period and later child psychopathology: A population cohort study. *J Am Acad Child Adolesc Psychiatry*. 2008;47(4):390-8. doi: 10.1097/CHI.0b013e31816429c2, PMID 18388761.
40. Wee KY, Skouteris H, Pier C, Richardson B, Milgrom J. Correlates of ante- and postnatal depression in fathers: A systematic review. *J Affect Disord*. 2011;130(3):358-77. doi: 10.1016/j.jad.2010.06.019, PMID 20599275.
41. Upadhyay RP, Chowdhury R, Aslyeh Salehi A, Sarkar K, Singh SK, Sinha B, Pawar A, Rajalakshmi AK, Kumar A. Postpartum depression in India: A systematic review and meta-analysis. *Bull World Health Organ*. 2017;95(10):706-717C. doi: 10.2471/BLT.17.192237, PMID 29147043.